



ORIGINAL

Assessing empathy among medical students: A comparative analysis using two different scales in a Brazilian medical school



Graziela Moreto^{a,*}, Itamar S. Santos^b, Pablo González Blasco^a, Leocir Pessini^c,
Paulo A. Lotufo^b

^a SOBRAMFA – Medical Education & Humanism, São Paulo, Brazil

^b Center for Clinical and Epidemiologic Research at Hospital Universitário, University of Sao Paulo, Sao Paulo, Brazil

^c São Camilo University Center, São Paulo, Brazil

Received 3 May 2017; accepted 14 July 2017

KEYWORDS

Empathy;
Medical education;
Humanism

Abstract

Purpose: The authors aimed to compare two different empathy scales across the six-year medical course, among undergraduate medical students attending a school of medicine that was established nine years ago in the Brazilian city of São Paulo.

Method: Cross-sectional study. The authors evaluated a sample of 296 students. The empathy evaluation was performed using the Jefferson Scale of Physician Empathy (JSPE), version for medical students, and Davis's multidimensional Interpersonal Reactivity Index (IRI-Brazilian version), applying both to each student simultaneously. The students were divided into three groups, according to their year on the medical course: Basic Group (1st and 2nd years), Clinical Group (3rd and 4th years) and Clerkship Group (5th and 6th years). Socio-demographic data and empathy scores, for both scales, were compared between these three groups using the Chi-square test and the Kruskal–Wallis test, as applicable.

Results: the JSPE scores were similar among the students from the Basic, Clinical and Clerkship groups ($p = 0.53$). On the other hand, the affective dimension of IRI-Brazilian version revealed a significantly lower score in the Clerkship Group ($p < 0.01$).

Conclusions: The authors suggest that the level of empathy can change, and in this case, the affective dimension was most affected during medical school.

© 2017 Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author.

E-mail address: graziela@sobramfa.com.br (G. Moreto).

PALABRAS CLAVE

Empatía;
Educación médica;
humanismo

Evaluación de la empatía entre los estudiantes de medicina: análisis comparativo utilizando dos escalas diferentes en una escuela de medicina de Brasil

Resumen

Objetivo: El objetivo de los autores fue comparar dos escalas diferentes de empatía a lo largo de la carrera de medicina de seis años, entre los estudiantes que asistían a una escuela de medicina fundada nueve años atrás en la ciudad brasileña de São Paulo.

Método: Estudio transversal. Los autores evaluaron una muestra de 296 estudiantes. La evaluación de la empatía se realizó utilizando la Escala de Empatía de los Médicos de Jefferson (JSPE), versión para estudiantes de medicina, y el Índice de Reactividad Interpersonal multidimensional de Davis (IRI-versión brasileña), aplicando ambos de manera simultánea a cada estudiante. Se dividió a los estudiantes en tres grupos, con arreglo a su curso de carrera de medicina: Grupo básico (1^{er} y 2^o años), Grupo Clínico (3^{er} y 4^o años) y Grupo de Prácticas (5^o y 6^o años). Se compararon los datos socio-demográficos y las puntuaciones sobre empatía entre estos tres grupos, utilizando las pruebas χ^2 y Kruskal-Wallis, según el caso.

Resultados: Las puntuaciones JSPE fueron similares entre los estudiantes de los grupos básico, clínico y de prácticas ($p=0,53$). Por otro lado, la dimensión afectiva de la versión brasileña de IRI reveló una puntuación inferior en el Grupo de Prácticas ($p < 0,01$).

Conclusiones: Los autores sugieren que el nivel de empatía puede cambiar y que, en este caso, la dimensión afectiva se vio más afectada durante la enseñanza de medicina.

© 2017 Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Some studies suggest that a drastic change takes place in the medical student during their training process. In the early years of medical training, most students are thrilled with the idea of becoming a physician, and are more sensitive to patients' suffering.^{1,2} During the medical course, this idealism can decrease, along with a process of 'dehumanization',³ in which the patient takes a secondary role in the medical practice.

One point that is acknowledged is that the medical student is exposed to a huge amount of scientific information that monopolizes their educational time,⁴ and rarely leaves room to address issues relating to the understanding of patients' expectations.⁵ To provide quality care, skills are needed to help the doctor not only to know about the disease, but also to understand the human being who is sick. Among these skills, empathy has proven to be crucial.⁶ Studies show that empathy among physicians is associated with better patient satisfaction,^{7,8} better adherence to treatment,^{8,9} more favorable clinical outcomes,¹⁰⁻¹³ and a decrease in the number of malpractice claims.¹⁴

The word empathy originates from the Greek word *empathēia*, meaning appreciation of the feelings of another person. The English term empathy was introduced by the psychologist Edward Titchener Bradner, and is used to mean the ability to understand another human being. Years later, Elmer Ernest Southard used the concept of empathy in the context of the doctor and patient, as a facilitator of diagnosis and therapy.¹⁵

There is disagreement among psychoanalysis scholars regarding the attributes involved in empathy. Some authors consider empathy to be a predominantly cognitive attribute,^{16,17} linking it to an understanding of the patient's

experiences and concerns, combined with the ability to communicate.¹⁸ Other authors see empathy essentially as belonging to the affective dimension, determined by the ability to experience the feelings of the other person.¹⁹ In this case, it is a consequence of the ability to be empathic, and it implies a spontaneous feeling of identification with the sufferer, in a process that involves emotion.

Some tools have recently been described to measure empathy. These include: the Interpersonal Reactivity Index (IRI) developed by Davis²⁰ with a version validated and adapted for Brazil (*Escala Multidimensional de Reatividade Interperssonal* – EMRI),²¹ the Hogan Empathy Scale²² and the Measure of Emotional Empathy developed by Mehrabian and Epstein.²³ The Jefferson Scale of Physician Empathy (JSPE) was specifically created for the patient care and medical education context.²⁴ Authors who have used some of these instruments have shown that during the training of different health professionals, empathy can provide positive or negative changes.⁶

A cross-sectional study by Chen et al.,²⁵ identified a low empathy score (measured through the JSPE scale) among 3rd year medical school students, compared to 2nd year students. A cohort study by Hojat et al.,⁶ identified an erosion of empathy (using JSPE as a tool to assess empathy) occurring in medical school, with the lowest scores among 3rd year students.

Understanding and assessing the degree of empathy of medical students during medical school is an important issue to be addressed during medical training. The aim of this study was to evaluate the degree of empathy among medical students in different years of the course at the Private University of São Paulo, using two different instruments: the JSPE (to assess the principally cognitive dimension of empathy for specific medical students within the medical

context) and the IRI-Brazilian version (to assess the cognitive and affective dimension for the population in general, not specific to the medical context).

Methodology

Study design

We included medical students from a private University in the city of São Paulo, which was established nine years ago. The total number of medical students in that University was 577. The students were invited to take part during the curricular classes. Adherence to the study was voluntary, and was related to the presence of the student in the classroom when the survey was administered. The data collection was carried out from September to December 2013. All the participants signed an informed consent form, and the study protocol was approved by the local ethics committee and by the ethics committee of the University of São Paulo School of Medicine.

Instruments

Two tools were used to evaluate the students' empathy: the Jefferson Scale of Physician Empathy (JSPE) and the IRI-Brazilian version. These were applied simultaneously to each participating student.

The choice of JSPE was related to its specificity for the medical student in a medical context, and the IRI-Brazilian version due to its multidimensional component, which assesses both the affective and cognitive dimensions of empathy.


The Jefferson Scale of Physician Empathy (JSPE) evaluates the cognitive dimension of empathy. It consists of 20 questions that are answered using the Likert scale of 7 points (1 = strongly disagree, 7 = strongly agree). The scores obtained for each question will be added together, resulting in a final score for empathy. Questions 1, 3, 6, 7, 8, 11, 12, 14, 18 and 19 have a reverse score.

The IRI-Brazilian version developed by Davis consists of 21 items that assess the cognitive and affective dimensions of empathy. The answer is given on a 5-point Likert scale (from 1 = "doesn't describe me very well" to 5 = "describes me very well"). The answers to questions 2, 3, 9, 10, 11 and 13 have a reverse score. The values obtained for each question will be added to the final score and dimension (cognitive and affective).

An isolated analysis was carried out of the mean Jefferson Scale scores, IRI-Brazilian version and subscales affective and cognitive dimension. For this analysis, the scores of each scale, from zero to 100 points were standardized and then compared to the average. This standardization was performed using the following formula:

$$\text{Score obtained} - \frac{\text{Minimum score}}{\text{Maximum score}} \\ - \text{Minimum score} \times 100$$

- Study variables From each participant, we obtained socio-demographic data: gender, year/semester of medical school, and the field of specialization the student was



Graphic 1 Analysis the field of specialization the student was interested in according to the year of graduation.

interested in. The specializations were classified in groups ("Clinic", "Surgery" and "Other")

- Statistical analysis The students were divided into 3 groups according to their year of graduation: "Basic" (1st and 2nd years), "Clinical" (3rd and 4th years) "Clerkship" (5th and 6th years). The socio-demographic data and empathy scores of both scales were compared between these 3 groups using the Chi-square test and the Kruskal-Wallis test, where applicable. We performed multiple linear regression models to assess the association between empathy scores according to year group (basic, clinical and clerkship). The models presented are crude and are adjusted for gender and intended specialty.

Results

Of the total of 577 medical students at the University studied, 296 (51%) participated in this study. [Graphic 1](#) shows the field of specialization the student was interested in according to the year of graduation (Basic – 1st and 2nd years, Clinical – 3rd and 4th years and Clerkship – 5th and 6th years).

Comparing the standardized averages of the two scales scores, we identified a significant difference between them, with the lowest average being obtained for the affective dimension of empathy. [Table 1](#) shows the maximum and minimum values of the scores obtained on each scale, as well as the standardized average.

Analyzing the empathy score, we found a significant decrease in empathy score in the affective dimension of IRI-Brazilian version in the Clerkship group ($p = 0.016$) ([Table 2](#)). The scores for the JSPE, IRI-Brazilian version and IRI-Brazilian version cognitive dimension showed no significant changes in the scores among the groups.

Comparing the empathy scores according to gender (men and women), we identified that women have higher empathy scores than men ($p < 0.01$ for IRI-Brazilian version affective dimension, IRI-Brazilian version and JSPE scores), but not for the cognitive component of IRI-Brazilian version scale ($p = 0.94$) ([Table 2](#)).

In the bivariate analysis, the JSPE, IRI-Brazilian version, and IRI-Brazilian version affective dimension scores were higher for students wishing to specialize in clinical specialties. In terms of intended specialty, there was no significant difference when using the cognitive dimension of the IRI-Brazilian version ([Table 2](#)).

Table 1 Scores of empathy scales and standardized mean.

	JSPE	IRI-Brazilian version	IRI-Brazilian version cognitive dimension	IRI-Brazilian version affective dimension
Minimum–maximum score	20–140	21–105	7–35	14–70
Reached maximum score	137	98	35	65
Reached minimum score	71	37	11	25
Mean	114	70	25	45
Standardized mean	78	58	64	55
P25–P75 standardized	72–86	52–65	53–75	46–64

Tables 3–6 show the results of the linear regression models. In the adjusted models, the JSPE scores were not significantly associated with the year of medical course or intended specialty. Men had lower empathy scores compared to women in the adjusted models. Analyzing the IRI-Brazilian version scores, we found lower scores in the clerkship group in the crude models, but this difference vanished after adjustment for gender. The full models showed lower scores in men and in those who intended to follow surgical careers. Analyzing the IRI-Brazilian version affective dimension score, the clerkship group, men, and those who intended to follow surgical careers were associated with lower scores in the full-adjusted models. The IRI-Brazilian version cognitive dimension scores showed no significant differences according to year of graduation, gender or intended specialization.

Discussion

We found that women obtained better scores for empathy compared to men, in both scales. Only in IRI-Brazilian version cognitive dimension was there no difference between genders. These findings related to gender were the same as identified in studies by Hojat et al.,⁶ Davis,²⁰ Eisenberg et al.,²⁶ Hasan et al.,²⁷ Mandel et al.,²⁸ Hojat et al.²⁹

About specialty, students intending to follow clinical careers had higher empathy scores, compared to those intending to follow a surgical career. The same results were found in studies by Hojat et al.,⁶ Coutts-van Dijk et al.,³⁰ Bailey.³¹

We would like to focus our discussion on another important finding, which was the possible affective erosion among students during medical school.

The results of this study demonstrate an important and challenging point in medical education, which is the medical student's affection. In all the statistical analyses, lower scores were obtained in the Clerkship group for the affective dimension of empathy assessed by the IRI-Brazilian version, suggesting that the affective dimension is the most affected during medical school. When we compared the scores of the JSPE and IRI-Brazilian version, the lowest score was obtained for the affective dimension of the IRI-Brazilian version, highlighting the importance of discussion on the subject of education in affection during medical school.

The universe of affection, feelings and emotions has had a leading role in the educational environment. The

classical pedagogical approach usually divides the educational objectives into three broad categories: cognitive, psychodynamic, and the education of affection. There is no way to objectively measure the growth or correct affective dynamic guidance of the student. In this context, it is more difficult to measure the danger of being forgotten, or placed under the purely arbitrary. Most of the time, the assessment of affective education is not taken into account when establishing educational objectives.³²

Educating emotions implies a serious process, which should initially be identified and then make them explicit and able to be worked by the teacher aiming to be used properly to promote effective patient doctor relationship. This educational process of educating affection requires tact, skill, avoid precipitation, promote learning to respect in any way, the physiological rhythm of emotion. You cannot force anyone to feel what one does not feel. What one can do is wait for a time of reflection over what one may feel, and from there develop affection, which is a true education process.³³

The emotions play an important role in the lives of medical students. They have to deal not only with their own emotions, but also with those of their patients. However, few components of their formal graduation are focused on their emotional lives.³⁴

A fact that supports this finding occurs when the student is exposed to medical practice during the clerkship (5th to 6th years). During this period, there is a predominant culture of keeping a certain distance from the patient, and avoiding emotional involvement. This behavior can be seen as a defense mechanism, a desire to distance oneself from the suffering of others, or a desire to avoid a sense of powerlessness over the situation.³⁵ When professionals, whether students or doctors, wrongly seek to protect their emotions under a cloak of impersonality, empathy is compromised.

It is understandable that without adequate preparation, medical students may express inappropriate emotional responses to situations commonly encountered in medical practice, such as pain, suffering and death – situations that make them aware of their own vulnerability.³⁶ When the student does not have the opportunity to work through their emotions, and to reflect and to share the feelings generated by them, they may develop an emotional detachment, seeking support based purely on their biomedical skills. But training in biomedical knowledge is given too much emphasis, merely drawing students further away from the humanist and ethical behavior that is always preferable.³⁷

Table 2 Compare empathy score according to gender, year of graduation, and intended specialization.

		Gender			Year of graduation				Intended specialization			
		Women	Men	<i>p</i>	Basic	Clinical	Clerkship	<i>p</i>	Clinic	Surgery	Others	<i>p</i>
JSPE	Median	118	115	0.004	115.5	116	118	0.53	119	115	108	0.05
	P25-P75	109-125	103-122		107-124	110-126.5	106.7-123.2		109.5-124	106-124	92.7-120.5	
	Mean	116.3	111.1		114.2	116	114		116.2	113.8	104	
	Standard deviation	11.0	13.8		11.6	12.2	12.2		11.3	12.5	19.77	
IRI-Brazilian version	Median	72	69	<0.001	71	71	70	0.17	72.5	70	66	0.007
	P25-P75	67-77	62-73.2		65.7-76	65.2-77.7	64-74		67.7-77	63.5-75	62-72	
	Mean	71.7	66.9		70.9	71.2	68.4		72	68.6	66.5	
	Standard deviation	8.6	9.8		8.8	10.2	9.2		8.6	9.8	6.9	
IRI-Brazilian version (cognitive)	Median	25	26	0.94	26	25	25	0.75	26	25	24	0.84
	P25-P75	23-28	22-29		23-29	22-28	22-28		23-28	22-29	23-29	
	Mean	25.2	25		25.2	24.8	25.3		25.3	25	24.8	
	Standard deviation	4.6	5.2		4.9	4.8	4.6		4.7	4.9	6.5	
IRI-Brazilian version (affective)	Median	46	42	<0.001	45	47	43	0.016	47	44	40	<0.001
	P25-P75	42-51.7	38-47		40.2-50	42-51.5	38-48		42-52	38-49	38-47	
	Mean	46.5	42		45.7	46.4	43.2		46.8	43.6	41.6	
	Standard deviation	6.8	7.3		6.7	8.0	7.2		6.6	7.8	5.8	

Table 3 Beta coefficients (95% confidence interval) associated with the score obtained in the JSPE empathy scale.

	Crude model	Model 1	Model 2
<i>Year</i>			
Basic	Reference	Reference	Reference
Clinical	+1.88 (−1.84 to 5.60)	+1.61 (−2.03 to 5.26)	+1.88 (−1.90 to 5.68)
Clerkship	+0.73 (−2.44 to 3.91)	+0.84 (−2.26 to 3.95)	+1.17 (−2.04 to 4.39)
<i>Gender</i>			
Women		Reference	Reference
Men		−5.29 (−8.19 to −2.39)	−4.32 (−7.27 to −1.37)
<i>Specialization</i>			
Clinic			Reference
Surgery			−2.48 (−5.37 to 0.41)
Others			−6.74 (−14.66 to 1.17)

The independent variables in adjusted model 1 are year (basic, clinical and clerkship) and gender.

The independent variables in adjusted model 2 are year (basic, clinical and clerkship), gender and intended specialization.

Table 4 Beta coefficients (95% confidence interval) associated with the score obtained in the IRI-Brazilian version.

	Crude model	Model 1	Model 2
<i>Year</i>			
Basic	Reference	Reference	Reference
Clinical	+0.25 (−2.65 to 3.16)	+0.19 (−2.63 to 3.02)	+0.14 (−2.82 to 3.12)
Clerkship	−2.50 (−5.00 to −0.02)	−2.25 (−4.68 to 0.17)	−2.10 (−4.62 to 0.40)
<i>Gender</i>			
Women		Reference	Reference
Men		−4.58 (−6.84 to −2.32)	−4.23 (−6.56 to −1.91)
<i>Specialization</i>			
Clinic			Reference
Surgery			−2.54 (−4.80 to −0.28)
Others			−4.25 (−10.39 to 1.89)

The independent variables in adjusted model 1 are year (basic, clinical and clerkship) and gender.

The independent variables in adjusted model 2 are year (basic, clinical and clerkship), gender and intended specialization.

Table 5 Beta coefficients (95% confidence interval) associated with the score obtained in the IRI-Brazilian version cognitive dimension.

	Crude model	Model 1	Model 2
<i>Year</i>			
Basic	Reference	Reference	Reference
Clinical	−0.43 (−1.93 to 1.07)	−0.43 (−1.94 to 1.07)	−0.61 (−2.23 to 0.99)
Clerkship	+0.06 (−1.22 to 1.34)	+0.06 (−1.22 to 1.35)	−0.02 (−1.37 to 1.33)
<i>Gender</i>			
Women		Reference	Reference
Men		−0.15 (−1.35 to 1.04)	−0.12 (−1.37 to 1.11)
<i>Specialization</i>			
Clinic			Reference
Surgery			−0.18 (−1.40 to 1.03)
Others			−0.31 (−3.66 to 3.02)

The independent variables in adjusted model 1 are year (basic, clinical and clerkship) and gender.

The independent variables in adjusted model 2 are year (basic, clinical and clerkship), gender, and intended specialization.

Table 6 Beta coefficients (95% confidence interval) associated with the score obtained in the IRI-Brazilian version affective dimensions.

	Crude model	Model 1	Model 2
<i>Year</i>			
Basic	Reference	Reference	Reference
Clinical	+0.67 (−1.50 to 2.93)	+0.60 (−1.55 to 2.77)	+0.75 (−1.52 to 3.03)
Clerkship	−2.48 (−4.41 to −0.54)	−2.25 (−4.11 to −0.39)	−1.99 (−3.91 to −0.07)
<i>Gender</i>			
Women		Reference	Reference
Men		−4.28 (−6.01 to −2.55)	−3.92 (−5.70 to −2.15)
<i>Specialization</i>			
Clinic			Reference
Surgery			−2.51 (−4.24 to −0.78)
Others			−4.08 (−8.80 to 0.62)

The independent variables in adjusted model 1 are year (basic, clinical and clerkship) and gender.

The independent variables in adjusted model 2 are year (basic, clinical and clerkship), gender and intended specialization.

To be empathic, it is necessary to see and understand the patient's suffering, and also to understand one's own reactions to the patient's suffering.³⁸ The exercise of self-critical reflection on the encounter with the patient, together with an understanding of one's own emotions, is a prerequisite to understanding the feelings of others, and is therefore an indicator of empathic ability.³⁹

Intervention possibilities

Bayne⁴⁰ highlights the importance of teacher education in this model of empathy process. It is essential for the development of empathy that the student experience empathic relationships with their teachers. This way, the student is more likely to develop an empathetic relationship with their patients.

Borrell⁴¹ strengthens the role of the teacher, stating that learning takes place through the "spread" of ethical attitudes from the teacher, demonstrated during their daily clinical practice.

There is a close relationship between life experiences and the development of empathy. Hence, the importance of learning by example, in an educational learning setting where medical students and young people are inspired by the attitudes of their teachers during practice with the patient. This model, which English-speaking educators call "tag-along", allows students to incorporate attitudes and behaviors in their approach to real patients, and to identify useful questions for their professional careers.⁴²

Beyond the example of the teacher, teaching and promoting empathy has the possibilities arising from humanistic and cultural training. The art that imitates life, offers diverse needs where the student is able to incorporate the features "hard to teach" (by traditional methods). Hence some authors emphasize the importance of art, literature and reflection in the student's own life, for the development of empathy.³⁶

The main goal of incorporating the humanities into medical education is to awaken attitudes and values that are often unexpected in the students themselves, who rely on

the scale of values, education and maturity that each possesses. The objectives are measured not so much by the final results, as the enlarged capacity of understanding of the human being that the process gives them, including what are called latent results.

Music has been shown to be a useful tool for enabling medical students to talk about themselves, ask questions, and discuss their expectations as a future doctor. It also helps revive ideals, dreams and life values.⁴³ The education of affectivity using film has also shown positive results in stimulating reflection among medical students.⁴⁴ Reflection is critical for the improvement of attitudes and behavior, as well as the incorporation of virtues and values.

Other instruments have also shown promising results in addressing issues related to the student's emotions, such as the use of literature, narratives⁴⁵ and opera.⁴⁶

Strengths and limitations of the study

This study is the application of consolidated fortress scales in the literature, which allows comparability with other studies. It also has some limitations, such as its single-center design, which affect the generalization and cross-sectional analysis. This enables the study of putative associations but impairs causal inference. Nevertheless, our findings of a trend toward lower empathy scores in the final years of graduation (statistically significant in IRI-Brazilian version affective dimension score) is consistent with the results of a longitudinal study by Hojat et al.,⁶ which identified an erosion of empathy during the medical course.

Conclusion

Our research suggests that the level of empathy can change and in this case, the affective dimension was the most affected during medical school. The identification of affective erosion strengthens the need to develop effective educational strategies that contribute to a good training medical for the professional future.

Conflict of interest

The authors of this article declare no conflict of interest.

References

1. Woloschuk W, Harasym PH, Temple W. Attitude change during medical school: a cohort study. *Med Educ.* 2004;38:522–34.
2. Mader EM, Roseamelia C, Morley CP. The temporal decline of idealism in two cohorts of medical students at one institution. *BMC Med Educ.* 2014;14:58.
3. Smith JK, Weaver DB. Capturing medical students' idealism. *Ann Fam Med.* 2006;4 Suppl. 1:58–60. S32–7.
4. Shapiro J. Does medical education promote professional alexithymia? A call for attending to the emotions of patients and self in medical training. *J Acad Med.* 2011;86:326–32.
5. Riess H, Kelley JM, Bailey RW, Dunn EJ, Phillips M. Empathy training for resident physicians: a randomized controlled trial of a neuroscience-informed curriculum. *J Gen Intern Med.* 2012.
6. Hojat M, Vergare MJ, Maxwell K, Brainard G, Herrine SK, Isenberg GA, et al. The devil is in the third year: a longitudinal study of erosion of empathy in medical school. *Acad Med.* 2009;84:1182–91.
7. Zachariae R, Pedersen CG, Jensen AB, Ehrnrooth E, Rossen PB, von der Maase H. Association of perceived physician communication style with patient satisfaction, distress, cancer-related self-efficacy, and perceived control over the disease. *Br J Cancer.* 2003;88:658–65.
8. Kim SS, Kaplowitz S, Johnston MV. The effects of physician empathy on patient satisfaction and compliance. *Eval Health Prof.* 2004;27:237–51.
9. Vermeire E, Hearnshaw H, Van Royen P, Denekens J. Patient adherence to treatment: three decades of research. A comprehensive review. *J Clin Pharm Ther.* 2001;26:331–42.
10. Hojat M, Louis DZ, Markham FW, Wender R, Rabinowitz C, Gonnella JS. Physicians' empathy and clinical outcomes for diabetic patients. *Acad Med.* 2011;86:359–64.
11. Di Blasi Z, Harkness E, Ernst E, Georgiou A, Kleijnen J. Influence of context effects on health outcomes: a systematic review. *Lancet.* 2001;357:757–62.
12. Rakel D, Barrett B, Zhang Z, Hoeft T, Chewning B, Marchand L, et al. Perception of empathy in the therapeutic encounter: effects on the common cold. *Patient Educ Couns.* 2011;85:390–7.
13. Rakel DP, Hoeft TJ, Barrett BP, Chewning BA, Craig BM, Niu M. Practitioner empathy and the duration of the common cold. *Fam Med.* 2009;41:494–501.
14. Levinson W, Roter DL, Mullooly JP, Dull VT, Frankel RM. Physician–patient communication: the relationship with malpractice claims among primary care physicians and surgeons. *JAMA.* 1997;277:553–9.
15. Hojat M. Empathy in patient care. In: *Antecedents, development, measurement and outcomes.* New York, NY: Springer; 2007.
16. Yamamoto S. Primate empathy: three factors and their combinations for empathy-related phenomena. *Wiley Interdiscip Rev Cogn Sci.* 2016.
17. Basch MF. Empathic understanding: a review of the concept and some theoretical considerations. *J Am Psychoanal Assoc.* 1983;31:101–26.
18. Hojat M, Gonnella JS, Nasca TJ, Mangione S, Vergare M, Magee M. Physician empathy: definition, components, measurement, and relationship to gender and specialty. *Am J Psychiatry.* 2002;159:1563–9.
19. Neumann M, Bensing J, Mercer S, Ernstmann N, Ommen N, Pfaff H. Analyzing the “nature” and “specific effectiveness” of clinical empathy: a theoretical overview and contribution towards a theory-based research agenda. *Patient Educ Counsel.* 2009;74:339–46.
20. Davis MH. Measuring individual differences in empathy: evidence for multidimensional approach. *J Pers Soc Psychol.* 1983;44:113–26.
21. Koller SH, Camino C, Ribeiro J. Adaptação e validação de duas escalas de empatia para uso no Brasil. *Estudos de psicologia (Campinas).* 2002;18:43–53.
22. Froman RD1, Peloquin SM. Rethinking the use of the Hogan empathy scale: a critical psychometric analysis. *Am J Occup Ther.* 2001;55:566–72.
23. Mehrabian A, Epstein NA. A measure of emotional empathy. *J Pers.* 1972;40:525–43.
24. Hojat M, Mangione S, Nasca TJ, Cohen MJM, Gonnella JS, Erdmann JB, et al. The Jefferson scale of physician empathy: development and preliminary psychometric data. *Educ Psychol Meas.* 2001;61:349–65.
25. Chen D, Lew R, Hershman W, Orlander J. A cross-sectional measurement of medical student empathy. *J Gen Intern Med.* 2007;22:1424–38.
26. Eisenberg N, Lennon R. Sex differences in empathy and related capacities. *Psychol Bull.* 1983;94:100–31.
27. Hasan S, Al-Sharqawi N, Dashti F, AbdulAziz M, Abdullah A, Shukkur M, et al. Level of empathy among medical students in Kuwait University Kuwait. *Med Princ Pract.* 2013;22:385–9.
28. Mandel ED, Schweinle WE. A study of empathy decline in physician assistant students at completion of first didactic year. *J Phys Assist Educ.* 2012;23:16–24.
29. Hojat M, Gonnella JS, Mangione S, Nasca TJ, Veloski JJ, Erdmann JB, et al. Empathy in medical students as related to academic performance, clinical competence and gender. *Med Educ.* 2002;36:522–7.
30. Coutts-van Dijk LC, Bray JH, Moore S, Rogers J. Prospective study of how students' humanism and psychosocial beliefs relate to specialty matching. *Acad Med.* 1997;72:1106–8.
31. Bailey BA. Empathy in medical students: assessment and relationship to specialty choice. *Diss Abstr Int.* 2001;62(6A):2024.
32. Blasco PG. Educação da afetividade através do cinema. Curitiba, PR, Brasil: IEF, Instituto de Ensino e Fomento; 2006.
- [33] Marias J. La educación sentimental. Madrid: Alianza Editorial; 1992.
34. Shapiro J. Does medical education promote professional alexithymia? A call for attending to the emotions of patients and self in medical training. *Acad Med.* 2011;86:326–32.
35. Lara JRL, Moral RR, Campayo JG. Por que algunos médicos se vuelven poco éticos (malvados?) com sus pacientes? *Atención Primaria.* 2009;41:646–9.
36. Larson EB, Yao X. Clinical empathy as emotional labor in patient–physician relationship. *JAMA.* 2005;293:1100–6.
37. Moreto G, González-Blasco P, Craice-De Benedetto MA. Reflexiones sobre la enseñanza de la empatía y la educación médica. *Aten Fam.* 2014;21:94–7.
38. Epstein RM. Mindful practice. *JAMA.* 1999;282:833–9.
39. Roter DL, Frankel RM, Hall JA, Sluyter D. *J Gen Intern Med.* 2006;21:28–34.
40. Bayne HB. Training medical students in empathic communication. *J Spec Group Work.* 2011;36:316–29.
41. Borrel-Carrio F, Epstein RM, Alenta HP. Profesionalidad y professionalism: fundamentos, contenidos, praxis y docencia. *Med Clin.* 2006;127:337–42.
42. Blasco PG, Roncoletta AFT, Moreto G, Levites MR, Janaudis MA. Accompanying physicians in their family practice: a primary care model for medical students' learning in Brazil. *Fam Med.* 2006;38:619–21.

43. Janaudis MA, Blasco PG, Ângelo M, Lotufo PA. Nos bailes da vida: a música facilitando a reflexão na educação médica. *Revista Brasileira de Medicina*. 2011;68:7-14.
44. Blasco PG. *Medicina de família & cinema: recursos humanísticos na educação médica*. São Paulo: Casa do Psicólogo; 2002.
45. Benedetto MAC. Entre dois continentes: literatura e narrativas humanizando médicos e pacientes. *O Mundo da Saúde*. 2010;34:311-9.
46. Blasco PG, Moreto G, Levites MR. Teaching humanities through opera: leading medical students to reflective attitudes. *Fam Med*. 2005;37:18-20.